

# MULTI-HYPERSUBSTITUTIONS AND COLORED SOLID VARIETIES

K. DENECKE, J. KOPPITZ, SL. SHTRAKOV

**ABSTRACT.** Hypersubstitutions are mappings which map operation symbols to terms. Terms can be visualized by trees. Hypersubstitutions can be extended to mappings defined on sets of trees. The nodes of the trees, describing terms, are labelled by operation symbols and by colors, i.e. certain positive integers. We are interested in mappings which map differently colored operation symbols to different terms. In this paper we extend the theory of hypersubstitutions and solid varieties to multi-hypersubstitutions and colored solid varieties. We develop the interconnections between such colored terms and multi-hypersubstitutions and the equational theory of Universal Algebra. The collection of all varieties of a given type forms a complete lattice which is very complex and difficult to study; multi-hypersubstitutions and colored solid varieties offer a new method to study complete sublattices of this lattice.

## 1. INTRODUCTION

Let  $X = \{x_1, \dots, x_n, \dots\}$  be a countably infinite set of variables, let  $X_n = \{x_1, \dots, x_n\}$  be a finite set and let  $(f_i)_{i \in I}$  be a set of operation symbols where  $f_i$  is  $n_i$ -ary. The sequence  $\tau := (n_i)_{i \in I}$  is called a type. In the usual way from variables and operation symbols we build up the set  $W_\tau(X)$  of all terms of type  $\tau$ . An algebra  $\mathcal{A} = (A; (f_i^{\mathcal{A}})_{i \in I})$  of type  $\tau$  is a pair consisting of a set  $A$  and an indexed set of operations defined on  $A$ . We denote by  $Alg(\tau)$  the class of all algebras of type  $\tau$ . If  $s, t \in W_\tau(X)$ , then the pair  $s \approx t$  is called an identity in the algebra  $\mathcal{A}$ , if the term operations  $s^{\mathcal{A}}$  and  $t^{\mathcal{A}}$  induced by the terms  $s$  and  $t$  on the algebra  $\mathcal{A}$  are equal. In this case we write  $\mathcal{A} \models s \approx t$ . The binary relation  $\models \subseteq Alg(\tau) \times W_\tau(X)^2$  gives rise to a Galois connection  $(Id, Mod)$  between the power sets of  $Alg(\tau)$  and  $W_\tau(X)^2$ , where  $Id$  and  $Mod$  are defined for  $K \subseteq Alg(\tau)$  and  $\Sigma \subseteq W_\tau(X)^2$  by

$$IdK := \{s \approx t \mid \forall \mathcal{A} \in K (\mathcal{A} \models s \approx t)\},$$

---

2000 *Mathematics Subject Classification.* Primary: 08A15; Secondary: 08A25.

*Key words and phrases.* Coloration of terms, Multi-hypersubstitutions, Colored solid varieties.

$$Mod\Sigma := \{\mathcal{A} \mid \forall s \approx t \in \Sigma (\mathcal{A} \models s \approx t)\}.$$

As a Galois connection,  $(Id, Mod)$  has the properties:

$$\Sigma_1 \subseteq \Sigma_2 \Rightarrow Mod\Sigma_2 \subseteq Mod\Sigma_1, \quad K_1 \subseteq K_2 \Rightarrow IdK_2 \subseteq IdK_1,$$

$$\Sigma \subseteq IdMod\Sigma, \quad K \subseteq ModIdK.$$

From these properties of the Galois connection  $(Id, Mod)$  we obtain that the fix points with respect to the closure operators

$$IdMod : \mathcal{P}(W_\tau(X)^2) \rightarrow \mathcal{P}(W_\tau(X)^2)$$

and

$$ModId : \mathcal{P}(Alg(\tau)) \rightarrow \mathcal{P}(Alg(\tau))$$

form complete lattices

$$\mathcal{L}(\tau) := \{K \mid K \subseteq Alg(\tau) \text{ and } ModIdK = K\}$$

$$\mathcal{E}(\tau) := \{\Sigma \mid \Sigma \subseteq W_\tau(X)^2 \text{ and } IdMod\Sigma = \Sigma\}$$

of all varieties of type  $\tau$  and of all equational theories of type  $\tau$ . These lattices are dually isomorphic.

Our next goal is to introduce two new closure operators on our sets  $Alg(\tau)$  and  $W_\tau(X)^2$  which give us complete sublattices of our two lattices  $\mathcal{L}(\tau)$  and  $\mathcal{E}(\tau)$ . The new operators are based on the concept of hypersatisfaction of an identity by a variety. We begin with the definition of a hypersubstitution. A complete study of hypersubstitutions may be found in [2].

A hypersubstitution of type  $\tau$  is a map which associates to every operation symbol  $f_i$  a term  $\sigma(f_i)$  of type  $\tau$ , of the same arity as  $f_i$ . Any hypersubstitution  $\sigma$  can be uniquely extended to a map  $\hat{\sigma}$  on the set  $W_\tau(X)$  of all terms of type  $\tau$  as follows:

- (i) If  $t = x_j$  for some  $j \geq 1$ , then  $\hat{\sigma}[t] = x_j$ ;
- (ii) if  $t = f_i(t_1, \dots, t_{n_i})$  for some  $n_i$ -ary operation symbol  $f_i$  and some terms  $t_1, \dots, t_{n_i}$ , then  $\hat{\sigma}[t] = \sigma(f_i)(\hat{\sigma}[t_1], \dots, \hat{\sigma}[t_{n_i}])$ .

Here the right side of (ii) means the composition of the term  $\sigma(f_i)$  and the terms  $\hat{\sigma}[t_1], \dots, \hat{\sigma}[t_{n_i}]$ .

We can define a binary operation  $\circ_h$  on the set  $Hyp(\tau)$  of all hypersubstitutions of type  $\tau$ , by taking  $\sigma_1 \circ_h \sigma_2$  to be the hypersubstitution which maps each fundamental operation symbol  $f_i$  to the term  $\hat{\sigma}_1[\sigma_2(f_i)]$ . That is,

$$\sigma_1 \circ_h \sigma_2 := \hat{\sigma}_1 \circ \sigma_2,$$

where  $\circ$  denotes the ordinary composition of functions. The operation  $\circ_h$  is associative. The *identity hypersubstitution*  $\sigma_{id}$  which maps every  $f_i$  to  $f_i(x_1, \dots, x_{n_i})$  is an identity element for this operation. Then  $\mathcal{Hyp}(\tau) := (\mathcal{Hyp}(\tau); \circ_h, \sigma_{id})$  is a monoid.

**Definition 1.1.** Let  $\mathcal{M}$  be any submonoid of  $\mathcal{Hyp}(\tau)$ . An algebra  $\mathcal{A}$  is said to  $M$ -hypersatisfy an identity  $u \approx v$  if for every hypersubstitution  $\sigma \in M$ , the identity  $\hat{\sigma}[u] \approx \hat{\sigma}[v]$  holds in  $\mathcal{A}$ . In this case we say that the identity  $u \approx v$  is an  $M$ -hyperidentity of  $\mathcal{A}$  and we write  $\mathcal{A} \models_{M-hyp} u \approx v$ .

For  $M = \mathcal{Hyp}(\tau)$  we write  $\mathcal{A} \models_{hyp} u \approx v$ .

An identity is called an  $M$ -hyperidentity of a variety  $V$  if it holds as an  $M$ -hyperidentity in every algebra in  $V$ . A variety  $V$  is called  $M$ -solid if every identity of  $V$  is an  $M$ -hyperidentity of  $V$ . When  $M$  is the whole monoid  $\mathcal{Hyp}(\tau)$ , an  $M$ -hyperidentity is called a *hyperidentity*, and an  $M$ -solid variety is called a *solid* variety.

Let  $\mathcal{M}$  be any submonoid of  $\mathcal{Hyp}(\tau)$ . Since  $M$  contains the identity hypersubstitution, any  $M$ -hyperidentity of a variety  $V$  is an identity of  $V$ . This means that the relation of  $M$ -hypersatisfaction, defined between  $Alg(\tau)$  and  $W_\tau(X)^2$ , is a subrelation of the relation of satisfaction from which we induced our Galois-connection  $(Id, Mod)$ . The new Galois-connection induced by the relation of  $M$ -hypersatisfaction is called  $(H_M Mod, H_M Id)$  and is defined on classes  $K$  and sets  $\Sigma$  as follows:

$$\begin{aligned} H_M Id K &= \{s \approx t \in W_\tau(X)^2 \mid s \approx t \text{ is an } M\text{-hyperidentity of } \mathcal{A} \text{ for all } \mathcal{A} \in K\}, \\ H_M Mod \Sigma &= \{\mathcal{A} \in Alg(\tau) \mid \text{all identities in } \Sigma \text{ are } M\text{-hyperidentities of } \mathcal{A}\}. \end{aligned}$$

The Galois-closed classes of algebras under this connection are the  $M$ -solid varieties of type  $\tau$ , which form a complete sublattice of the lattice of all varieties of type  $\tau$ . Thus studying  $M$ -solid and solid varieties is a way to study complete sublattices of the lattice of all varieties of a given type.

We now introduce some closure operators on the two sets  $Alg(\tau)$  and  $W_\tau(X)^2$ . For equations we define

$$\chi_M^E[u \approx v] := \{\hat{\sigma}[u] \approx \hat{\sigma}[v] \mid \sigma \in M\}.$$

For any set  $\Sigma$  of identities we set

$$\chi_M^E[\Sigma] = \bigcup_{u \approx v \in \Sigma} \chi_M^E[u \approx v].$$

Hypersubstitutions can also be applied to algebras, as follows. Given an algebra  $\mathcal{A} = (A; (f_i^{\mathcal{A}})_{i \in I})$  and a hypersubstitution  $\sigma$ , we define the algebra  $\sigma(\mathcal{A}) = (A; (f_i^{\sigma(\mathcal{A})})_{i \in I}) = (A; (\sigma(f_i)^{\mathcal{A}})_{i \in I})$ . This algebra is called the *derived algebra* determined by  $\mathcal{A}$  and  $\sigma$ . Notice that by definition it is of the same type as the algebra  $\mathcal{A}$ . Now we define an operator  $\chi_M^A$  on the set  $\text{Alg}(\tau)$ , first on individual algebras and then on classes  $K$  of algebras, by

$$\chi_M^A[\mathcal{A}] = \{\sigma[\mathcal{A}] \mid \sigma \in M\},$$

and

$$\chi_M^A[K] = \bigcup_{\mathcal{A} \in K} \chi_M^A[\mathcal{A}].$$

If  $M = \text{Hyp}(\tau)$  the operators are denoted by  $\chi^A$  and  $\chi^E$ .

Let  $\tau$  be a fixed type and let  $\mathcal{M}$  be any submonoid of  $\text{Hyp}(\tau)$ . The two operators  $\chi_M^E$  and  $\chi_M^A$  are closure operators and are connected by the condition

$$\chi_M^A[\mathcal{A}] \text{ satisfies } u \approx v \text{ iff } \mathcal{A} \text{ satisfies } \chi_M^E[u \approx v].$$

The following propositions are also obvious (see [1] or [2]).

**Theorem 1.2.** *Let  $K \subseteq \text{Alg}(\tau)$  and  $\Sigma \subseteq W_\tau(X)$ . Then there holds*

- (i)  $H_M \text{Mod} \Sigma = \text{Mod} \chi_M^E[\Sigma],$
- (ii)  $H_M \text{Mod} \Sigma \subseteq \text{Mod} \Sigma,$
- (iii)  $\chi_M^A[H_M \text{Mod} \Sigma] = H_M \text{Mod} \Sigma,$
- (iv)  $\chi_M^E[\text{Id} H_M \text{Mod} \Sigma] = \text{Id} H_M \text{Mod} \Sigma,$
- (v)  $H_M \text{Mod} H_M \text{Id} K = \text{Mod} \text{Id} \chi_M^A[K],$
- (i')  $H_M \text{Id} K = \text{Id} \chi_M^A[K],$
- (ii')  $H_M \text{Id} K \subseteq \text{Id} K,$
- (iii')  $\chi_M^E[H_M \text{Id} K] = H_M \text{Id} K,$
- (iv')  $\chi_M^A[\text{Mod} H_M \text{Id} K] = \text{Mod} H_M \text{Id} K,$
- (v')  $H_M \text{Id} H_M \text{Mod} \Sigma = \text{Id} \text{Mod} \chi_M^E[\Sigma].$

$M$ -solid varieties can be characterized by the following theorem:

**Theorem 1.3.** *Let  $\mathcal{M}$  be a monoid of hypersubstitutions of type  $\tau$ . For any variety  $V$  of type  $\tau$ , the following conditions are equivalent:*

- (i)  $V = H_M \text{Mod} H_M \text{Id} V,$
- (ii)  $\chi_M^A[V] = V,$
- (iii)  $\text{Id} V = H_M \text{Id} V,$
- (iv)  $\chi_M^E[\text{Id} V] = \text{Id} V.$

And dually, for any equational theory  $\Sigma$  of type  $\tau$ , the following conditions are equivalent:

- (i')  $\Sigma = H_M \text{Id} H_M \text{Mod} \Sigma,$
- (ii')  $\chi_M^E[\Sigma] = \Sigma,$
- (iii')  $\text{Mod} \Sigma = H_M \text{Mod} \Sigma,$
- (iv')  $\chi_M^A[\text{Mod} \Sigma] = \text{Mod} \Sigma.$

■

A variety which satisfies the condition (i) is called an  $M$ -hyperequational class.

The subrelation  $\models_{M\text{-hyp}}$  of  $\models$  satisfies the following condition:

$$\forall K \subseteq \text{Alg}(\tau) \quad \forall \Sigma \subseteq W_\tau(X)^2 \quad ((H_M \text{Mod} \Sigma = K \wedge H_M \text{Id} K = \Sigma) \Rightarrow \text{Mod} \Sigma = K \wedge \text{Id} K = \Sigma).$$

Such subrelations are called Galois-closed. The Galois connections induced by Galois-closed subrelations of a given relation generate complete sublattices of the complete lattices generated by the Galois connection induced by that given relation [3]. Moreover we have the following result.

**Theorem 1.4.** *Let  $\mathcal{M}$  be a monoid of hypersubstitutions of type  $\tau$ . Then the class  $\mathcal{S}_M(\tau)$  of all  $M$ -solid varieties of type  $\tau$  forms a complete sublattice of the lattice  $\mathcal{L}(\tau)$  of all varieties of type  $\tau$ . Dually, the class of all  $M$ -hyperequational theories forms a complete sublattice of the lattice of all equational theories of type  $\tau$ . □*

When  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are both submonoids of  $\mathcal{Hyp}(\tau)$  and  $\mathcal{M}_1$  is a submonoid of  $\mathcal{M}_2$ , then the corresponding complete lattices satisfy  $\mathcal{S}_{\mathcal{M}_2}(\tau) \subseteq \mathcal{S}_{\mathcal{M}_1}(\tau)$ . As a special case, for any  $\mathcal{M} \subseteq \mathcal{Hyp}(\tau)$  we see that the lattice  $\mathcal{S}(\tau)$  of all solid varieties of type  $\tau$  is always a sublattice of the lattice  $\mathcal{S}_M(\tau)$ . At the other extreme, for the smallest possible submonoid  $\mathcal{M} = \{\sigma_{id}\}$  the corresponding lattice of  $M$ -solid varieties is the whole lattice  $\mathcal{L}(\tau)$  of all varieties of type  $\tau$ . Thus we obtain a range of complete sublattices of  $\mathcal{L}(\tau)$  to  $\mathcal{S}(\tau)$ .

Our aim is to transfer this theory to another kind of hypersubstitution and to colored terms.

## 2. MULTI-HYPERSUBSTITUTIONS AND COLORATIONS OF TERMS

In a term a certain operation symbol may occur more than once. To distinguish between the different occurrences of the same operation symbol we assign to each occurrence of any operation symbol a color. Representing a term by a tree, we get a vertex-colored graph. To apply different hypersubstitutions to the same operation symbol, if it is differently colored, we define the concept of a multi-hypersubstitution.

**Definition 2.1.** A map  $\rho$  from  $\mathbb{N}$  into  $Hyp(\tau)$  is called a multi- hypersubstitution. Let  $Hyp^{\mathbb{N}}$  be the set of all multi-hypersubstitutions.

To distinguish between different occurrences of the same operation symbol in a term  $t$  we assign to each operation symbol in  $t$  an address in  $t$ , i.e. an element of a given set. Usually addresses are sequences of natural numbers. Let  $add(t)$  be the set of all addresses of the term  $t$ . We introduce the concept of a coloration to allow that equal operation symbols at different places are considered differently. On the other hand, a coloration allows that equal operation symbols with different addresses are equally colored. A coloration of a type  $\tau$  is defined in the following way.

**Definition 2.2.** Any mapping  $\alpha_t$  from  $add(t)$ ,  $t \in W_{\tau}(X) \setminus X$ , into  $\mathbb{N}$  is called a coloration of the term  $t$ . We denote by  $C(t)$  the set of all colorations of the term  $t$ . A set  $C \subseteq \bigcup \{C(r) \mid r \in W_{\tau}(X)\}$  with  $|C(r) \cap C| = 1$  for all  $r \in W_{\tau}(X) \setminus X$  is called a coloration of  $W_{\tau}(X)$ .

Using colorations, multi- hypersubstitutions can be extended to mappings defined on terms. In a first step we extend multi- hypersubstitutions to mappings from the set  $Sub(t)$  of all subterms of a given term  $t$  to the set of terms.

**Definition 2.3.** Let  $C$  be a coloration of  $W_{\tau}(X)$ ,  $t \in W_{\tau}(X)$  with the coloration  $\alpha \in C$ ,  $s \in Sub(t)$ , and let  $\rho$  be a multi-hypersubstitution.

- (i) If  $s \in X$  then  $\widehat{\rho}_{C,t}[s] := s$ .
- (ii) If  $s = f_i(s_1, \dots, s_{n_i})$  with  $i \in I$  and  $s_1, \dots, s_{n_i} \in Sub(t)$ , where  $f_i$  has the address  $a$  in  $t$  then

$$\widehat{\rho}_{C,t}[s] := \rho(\alpha(a))(f_i)(\widehat{\rho}_{C,t}[s_1], \dots, \widehat{\rho}_{C,t}[s_{n_i}]),$$

Using the mappings  $\widehat{\rho}_{C,t}[t] : Sub(t) \rightarrow W_{\tau}(X)$  for terms  $t \in W_{\tau}(X)$  we can extend multi-hypersubstitutions to mappings defined on terms.

**Definition 2.4.** Let  $C$  be a coloration of  $W_\tau(X)$  and  $\alpha \in C$  be the coloration of a term  $t \in W_\tau(X)$ . Then for a multi-hypersubstitution  $\rho$  we put  $\widehat{\rho}_C[t] := \widehat{\rho}_{C,t}[t]$ .

The following example shows that the composition of two multi-hypersubstitutions does not be a multi-hypersubstitution. For this we consider the type  $\tau = (2)$ , the terms  $s = f(y, f(y, x))$  and  $t = f(f(x, y), y)$  with the following coloration  $C$ :

$$\begin{aligned} \alpha_t(a) &= 0 \text{ for all } a \in \text{add}(t); \\ \alpha_s(q) &= 0, \text{ where } s \text{ is the address of the leftmost } f \text{ in } s; \\ \alpha_s(a) &= 1 \text{ for all } a \in \text{add}(s), a \neq q. \end{aligned}$$

Let  $\rho \in \text{Hyp}(\tau)^\mathbb{N}$  be a multi-hypersubstitution with  $\rho(0) = \sigma_{yx}$  and  $\rho(a) = \sigma_{id}$  for  $a \in \mathbb{N} \setminus \{0\}$ . Then we have  $\widehat{\rho}_C[t] = s$  and  $\widehat{\rho}_C[t] = f(f(y, x), y)$ . Since all operation symbols in  $t$  have the same color, we can not find a multi-hypersubstitution which provides  $f(f(y, x), y)$  by application on  $t$ .

If all addresses of a term have the same color, then a multi-hypersubstitution can be replaced on that term by one of its components, i.e. by an ordinary hypersubstitution as the following lemma shows.

**Lemma 2.5.** *Let  $C$  be a coloration of  $W_\tau(X)$ ,  $\rho \in \text{Hyp}(\tau)^\mathbb{N}$ , and  $t \in W_\tau(X)$  such that there is an  $n \in \mathbb{N}$  with  $\alpha_t(a) = n$  for all  $a \in \text{add}(t)$ . Then  $\widehat{\rho}_C[t] = \widehat{\rho(n)}[t]$ .*

**Proof:** Since  $\widehat{\rho}_C[t] = \widehat{\rho}_{C,t}[t]$  we show by induction that  $\widehat{\rho(n)}[s] := \widehat{\rho}_{C,t}[s]$  for each  $s \in \text{Sub}(t)$ . If  $s \in X$  then  $\widehat{\rho}_{C,t}[s] = s = \widehat{\rho(n)}[s]$ .

Let  $s = f_i(s_1, \dots, s_{n_i})$  and suppose that  $\widehat{\rho}_{C,t}[s_j] = \widehat{\rho(n)}[s_j]$  for  $1 \leq j \leq n_i$  then

$$\begin{aligned} \widehat{\rho}_{C,t}[f_i(s_1, \dots, s_{n_i})] &= \rho(\alpha_t(a))(f_i)(\widehat{\rho}_{C,t}[s_1], \dots, \widehat{\rho}_{C,t}[s_{n_i}]) \\ &\quad (a \text{ denotes the address of } f_i \text{ in } t) \\ &= \rho(\alpha_t(a))(f_i)(\widehat{\rho(n)}[s_1], \dots, \widehat{\rho(n)}[s_{n_i}]) \\ &\quad (\text{by the hypothesis}) \\ &= \widehat{\rho(n)}(f_i)(\widehat{\rho(n)}[s_1], \dots, \widehat{\rho(n)}[s_{n_i}]) \\ &= \widehat{\rho(n)}[f_i(s_1, \dots, s_{n_i})]. \end{aligned}$$

■

## 3. COLORED SOLID VARIETIES

Using multi-hypersubstitutions we define operators corresponding to  $\chi_M^A, \chi_M^E$  of the introduction and apply these operators to sets of equations and to classes of algebras.

**Definition 3.1.** Let  $C$  be a coloration of  $W_\tau(X)$  and  $\Sigma \subseteq W_\tau(X)^2$ . Then we put  $\chi_C^e[\Sigma] := \{\widehat{\rho}_C[u] \approx \widehat{\rho}_C[v] \mid u \approx v \in \Sigma, \rho \in \text{Hyp}(\tau)^\mathbb{N}\} = \chi_C^{e,0}[\Sigma]$ . For  $n \in \mathbb{N}$ , we put  $\chi_C^{e,n+1}[\Sigma] := \chi_C^e[\chi_C^{e,n}[\Sigma]]$ . Let  $\chi_C^E[\Sigma] := \bigcup \{\chi_C^{e,n}[\Sigma] \mid n \in \mathbb{N}\}$ .

For any set  $\Sigma \subseteq W_\tau(X)^2$  we have  $\chi_C^e[\Sigma] = \Sigma$  iff  $\chi_C^E[\Sigma] = \Sigma$ . Indeed, If  $\chi_C^e[\Sigma] = \Sigma$  then it is easy to see that  $\chi_C^{e,n}[\Sigma] = \Sigma$  for all  $n \in \mathbb{N}$  and thus  $\chi_C^E[\Sigma] = \Sigma$ . Conversely, if  $\chi_C^E[\Sigma] = \Sigma$  then  $\chi_C^e[\Sigma] \subseteq \chi_C^E[\Sigma]$  provides  $\chi_C^e[\Sigma] \subseteq \Sigma$ . Since  $\rho_{id} \in \text{Hyp}(\tau)^\mathbb{N}$  we have  $\Sigma \subseteq \chi_C^e[\Sigma]$  and altogether,  $\chi_C^e[\Sigma] = \Sigma$ .

**Definition 3.2.** Let  $C$  be a coloration of  $W_\tau(X)$  and let  $V$  be a variety of type  $\tau$ .  $V$  is called  $C$ -colored solid if  $IdV = \chi_C^E[IdV]$ .

**Example 3.3.** We consider the following example of a  $C$ -colored variety of type  $\tau = (2)$ . Let  $RB = \text{Mod}\{x(yz) \approx (xy)z \approx xz, x^2 \approx x\}$  be the variety of rectangular bands. It is well-known that  $RB$  is solid (see e.g. [6]). The set  $IdRB$  of all identities satisfied in  $RB$  is the set of all equations  $s \approx t$  such that the first variable of  $s$  agrees with the first variable of  $t$  and the last variable of  $s$  agrees with the last variable of  $t$ . For  $t \in W_{(2)}(X)$  such that  $t$  starts and ends with the same variable we define  $\alpha_t : \text{add}(t) \rightarrow \mathbb{N}$  with  $\alpha_t(a) = 1$  for all  $a \in \text{add}(t)$  and if  $t$  starts and ends with different variables, then we define  $\alpha_t(a) = 2$  for all  $a \in \text{add}(t)$ . Let  $s \approx t \in IdRB$  and let  $\rho$  be a multi-hypersubstitution. Then it is easy to see that  $\widehat{\rho}_C[s] \approx \widehat{\rho}_C[t] \in IdRB$  using solidity.

**Definition 3.4.** Let  $C$  be a coloration of  $W_\tau(X)$ , let  $\rho \in \text{Hyp}(\tau)^\mathbb{N}$  and let  $\mathcal{A} = (A; (f_i^A)_{i \in I})$  be an algebra of type  $\tau$ . Then we define:  $\rho[\mathcal{A}] := (A; (f_i^{\rho[\mathcal{A}]})_{i \in I})$  where  $f_i^{\rho[\mathcal{A}]} = \widehat{\rho}_C[f_i(x_1, \dots, x_{n_i})]^\mathcal{A}$  for  $i \in I$ .

**Definition 3.5.** Let  $C$  be a coloration of  $W_\tau(X)$  and  $K$  be a class of algebras of type  $\tau$ . Then we put  $\chi_C^A[K] := \{\rho[\mathcal{A}] \mid \mathcal{A} \in K, \rho \in \text{Hyp}(\tau)^\mathbb{N}\}$ .

It is easy to check that  $\chi_C^E$  and  $\chi_C^A$  have the properties of a completely additive closure operator. From Lemma 3.6 below will follow that a  $C$ -colored solid variety is solid. Our new closure operators are connected with the operators defined in the Introduction.

**Lemma 3.6.** Let  $C$  be a coloration of  $W_\tau(X)$ . Then



- (i)  $\chi^E[\Sigma] \subseteq \chi_C^E[\Sigma]$  for each  $\Sigma \subseteq W_\tau(X)^2$ .
- (ii)  $\chi_C^A[K] = \chi^A[K]$  for each  $K \subseteq \text{Alg}(\tau)$ .

**Proof:** (i)  $\chi_C^E[\Sigma] \subseteq \chi_C^E[\Sigma]$  we have to show that  $\chi^E[\Sigma] \subseteq \chi_C^E[\Sigma]$ . Let  $\sigma \in \text{Hyp}(\tau)$ . Then we consider the multi-hypersubstitution  $\rho \in \text{Hyp}(\tau)^\mathbb{N}$  with  $\rho(a) = \sigma$  for all  $a \in \mathbb{N}$ . If  $t \in W_\tau(X)$  then we will check that  $\widehat{\sigma}[t] = \widehat{\rho}_C[t]$ . Since  $\widehat{\rho}_C[t] = \widehat{\rho}_{C,t}[t]$  we show that  $\widehat{\sigma}[s] := \widehat{\rho}_{C,t}[s]$  for each  $s \in \text{Sub}(t)$  by induction on the complexity of the term  $s$ .

If  $s \in X$  then  $\widehat{\sigma}[s] = s = \widehat{\rho}_{C,t}[s]$ .

Assume that  $s = f_i(s_1, \dots, s_{n_i})$  with  $i \in I$  and  $s_1, \dots, s_{n_i} \in \text{Sub}(t)$  and suppose inductively that  $\widehat{\sigma}[s_k] = \widehat{\rho}_{C,t}[s_k]$  for  $1 \leq k \leq n_i$ . Then for a suitable  $a \in \mathbb{N}$  we have

$$\begin{aligned} \widehat{\rho}_{C,t}[s] &= \rho(a)(f_i)(\widehat{\rho}_{C,t}[s_1], \dots, \widehat{\rho}_{C,t}[s_{n_i}]) \\ &= \sigma(f_i)(\widehat{\sigma}[s_1], \dots, \widehat{\sigma}[s_{n_i}]) = \widehat{\sigma}[f_i(s_1, \dots, s_{n_i})]. \end{aligned}$$

- (ii) Let  $\mathcal{A} \in K$ . For  $i \in I$  let  $a_i$  be the color of  $f_i$  in the fundamental term  $f_i(x_1, \dots, x_{n_i})$ .

Let  $\rho \in \text{Hyp}(\tau)^\mathbb{N}$ . Then we consider the hypersubstitution  $\sigma \in \text{Hyp}(\tau)$  with  $\sigma(f_i) = \rho(a_i)(f_i)$  for  $i \in I$  and have

$$\begin{aligned} f_i^{\rho[\mathcal{A}]} &= \widehat{\rho}_C[f_i(x_1, \dots, x_{n_i})]^\mathcal{A} = (\rho(a_i)(f_i)(x_1, \dots, x_{n_i}))^\mathcal{A} \\ &= \sigma(f_i)(x_1, \dots, x_{n_i})^\mathcal{A} = \sigma(f_i)^\mathcal{A}. \end{aligned}$$

This shows that  $\chi_C^A[\mathcal{A}] \subseteq \chi^A[\mathcal{A}]$ .

Let  $\sigma \in \text{Hyp}(\tau)$ . Then we consider the multi-hypersubstitution  $\rho \in \text{Hyp}(\tau)^\mathbb{N}$  with  $\rho(a) = \sigma$  for all  $a \in \mathbb{N}$ . Now we have

$$\begin{aligned} f_i^{\sigma[\mathcal{A}]} &= \sigma(f_i)^\mathcal{A} = (\sigma(f_i)(x_1, \dots, x_{n_i}))^\mathcal{A} = (\rho(a_i)(f_i)(x_1, \dots, x_{n_i}))^\mathcal{A} \\ &= \widehat{\rho}_C[f_i(x_1, \dots, x_{n_i})]^\mathcal{A} = f_i^{\rho[\mathcal{A}]}. \end{aligned}$$

This shows that  $\chi^A[\mathcal{A}] \subseteq \chi_C^A[\mathcal{A}]$ . Altogether we have  $\chi_C^A[\mathcal{A}] = \chi^A[\mathcal{A}]$  and thus  $\chi_C^A[K] = \chi^A[K]$ .  $\blacksquare$

Using the operators  $\chi_C^A$  and  $\chi_C^E$  we define two new relations between  $\text{Alg}(\tau)$  and  $W_\tau(X)^2$ .

**Definition 3.7.** Let  $C$  be a coloration of  $W_\tau(X)$ . Then we put

$$R_1 := \{(\mathcal{A}, s \approx t) \mid \mathcal{A} \in \text{Alg}(\tau), s \approx t \in W_\tau(X)^2, \chi_C^E[s \approx t] \subseteq \text{Id}\mathcal{A}\};$$

$$R_2 := \{(\mathcal{A}, s \approx t) \mid \mathcal{A} \in \text{Alg}(\tau), s \approx t \in W_\tau(X)^2, s \approx t \in \text{Id}\chi_C^A[\mathcal{A}]\};$$

$$CMod\Sigma := \{\mathcal{A} \mid \{\mathcal{A}\} \times \Sigma \subseteq R_1\} \text{ for } \Sigma \subseteq W_\tau(X)^2;$$

$\mathcal{C}IdK := \{s \approx t \mid K \times \{s \approx t\} \subseteq R_1\}$  for  $K \subseteq Alg(\tau)$ .

Because of Lemma 3.6 (ii) the relation  $R_2$  agrees with the relation  $\models_{hyp}$ .

Since  $\models_{hyp}$  is a Galois closed subrelation of  $\models$ , the relation  $R_2$  has the same property. Since we are more interested in  $R_1$ , we ask whether  $R_1$  is a Galois closed subrelation of  $\models$ . At first we have the following property:

**Proposition 3.8.** *Let  $C$  be a coloration of  $W_\tau(X)$ . Then  $R_1$  is a subrelation of  $\models$  such that for  $K \subseteq Alg(\tau)$  and all  $\Sigma \subseteq W_\tau(X)^2$  the following holds: If  $\Sigma = \mathcal{C}IdK$  and  $K = \mathcal{C}Mod\Sigma$  then  $K = Mod\Sigma$ .*

**Proof:** At first we show that  $R_1$  is a subrelation of  $\models$ . For this let  $(\mathcal{A}, s \approx t) \in R_1$ . Then  $\chi_C^E[s \approx t] \subseteq Id\mathcal{A}$  and  $s \approx t \in Id\mathcal{A}$  since  $\chi_C^E$  is a closure operator. Thus  $(\mathcal{A}, s \approx t) \in \models$ .

Now we show that  $\Sigma = \chi_C^E[\Sigma]$ . Since  $\chi_C^E$  is a closure operator, we have  $\Sigma \subseteq \chi_C^E[\Sigma]$ .

Conversely let  $u \approx v \in \chi_C^E[\Sigma]$ . Then there is an  $s \approx t \in \Sigma$  with  $u \approx v \in \chi_C^E[s \approx t]$ , i.e.  $\chi_C^E[u \approx v] \subseteq \chi_C^E[\chi_C^E[s \approx t]] \subseteq \chi_C^E[s \approx t]$ . From  $s \approx t \in \Sigma = \mathcal{C}IdK$  it follows  $\chi_C^E[s \approx t] \subseteq IdK$  and  $\chi_C^E[u \approx v] \subseteq \chi_C^E[s \approx t] \subseteq IdK$ , i.e.  $u \approx v \in \mathcal{C}IdK$ . This shows  $\chi_C^E[\Sigma] \subseteq \Sigma$ . Now we have

$$\begin{aligned} Mod\Sigma &= \{ \mathcal{A} \mid \mathcal{A} \in Alg(\tau), \Sigma \subseteq Id\mathcal{A} \} \\ &= \{ \mathcal{A} \mid \mathcal{A} \in Alg(\tau), \chi_C^E[\Sigma] \subseteq Id\mathcal{A} \} \\ &= \mathcal{C}Mod\Sigma \\ &= K. \end{aligned}$$

■

To obtain a characterization of colored solid varieties we check at first the conditions of Theorem 1.2 for M-solid varieties.

**Lemma 3.9.** *Let  $C$  be a coloration of  $W_\tau(X)$ , let  $\Sigma \subseteq W_\tau(X)^2$ , and let  $K$  be a class of algebras of type  $\tau$ . Then*

$$\chi_C^E[\Sigma] \subseteq IdK \Leftrightarrow \chi_C^E[\Sigma] \subseteq Id\chi_C^A[K].$$

**Proof:** Because of  $K \subseteq \chi_C^A[K] \Rightarrow Id\chi_C^A[K] \subseteq IdK$  we get  $\chi_C^E[\Sigma] \subseteq Id\chi_C^A[K] \Rightarrow \chi_C^E[\Sigma] \subseteq IdK$ .

By Lemma 3.6 (i) and the definition of the closure operators  $\chi_C^E$  and  $\chi_C^A$  we have  $\chi_C^E[\Sigma] = \chi_C^E[\chi_C^E[\Sigma]] \supseteq \chi_C^E[\chi_C^A[\Sigma]] \supseteq \chi_C^E[\Sigma]$ , i.e.  $\chi_C^E[\Sigma] = \chi_C^E[\chi_C^A[\Sigma]]$ .

Consequently we have

$\chi_C^E[\Sigma] \subseteq IdK \Rightarrow \chi^E[\chi_C^E[\Sigma]] \subseteq IdK \Rightarrow \chi_C^E[\Sigma] \subseteq Id\chi_C^A[K]$ ,  
since  $(\chi^E, \chi^A)$  is a conjugate pair. Then by Lemma 3.6,  $\chi_C^E[\Sigma] \subseteq Id\chi_C^A[K]$ . ■

As a consequence we have

$$\mathcal{C}Mod\Sigma = \{\mathcal{A} \mid \mathcal{A} \in Alg(\tau) \text{ and } \chi_C^E[\Sigma] \subseteq Id\chi_C^A[\mathcal{A}]\}. \quad (*)$$

Now we prove that the sets of the form  $\mathcal{C}IdK$  and  $\mathcal{C}Mod\Sigma$  are closed under the operators  $\chi_C^E$  and  $\chi_C^A$ , respectively.

**Lemma 3.10.** *Let  $C$  be a coloration of  $W_\tau(X)$ .*

- (i) *For  $K \subseteq Alg(\tau)$  there holds  $\chi_C^E[\mathcal{C}IdK] = \mathcal{C}IdK$ .*
- (ii) *For  $\Sigma \subseteq W_\tau(X)^2$  there holds  $\chi_C^A[\mathcal{C}Mod\Sigma] = \mathcal{C}Mod\Sigma$ .*

**Proof:** (i) Clearly,  $\mathcal{C}IdK \subseteq \chi_C^E[\mathcal{C}IdK]$ . Let  $u \approx v \in \chi_C^E[\mathcal{C}IdK]$ . Then there is an equation  $s \approx t \in \mathcal{C}IdK$  with  $u \approx v \in \chi_C^E[s \approx t]$ . Since  $\chi_C^E$  is a closure operator we have  $\chi_C^E[u \approx v] \subseteq \chi_C^E[\chi_C^E[s \approx t]] = \chi_C^E[s \approx t]$ . From  $s \approx t \in \mathcal{C}IdK$  it follows  $\chi_C^E[s \approx t] \subseteq IdK$ . Then  $\chi_C^E[u \approx v] \subseteq IdK$ , thus  $u \approx v \in \mathcal{C}IdK$ .

(ii) Clearly,  $\mathcal{C}Mod\Sigma \subseteq \chi_C^A[\mathcal{C}Mod\Sigma]$ . Let  $\mathcal{A} \in \chi_C^A[\mathcal{C}Mod\Sigma]$ . Then there is a  $\mathcal{B} \in \mathcal{C}Mod\Sigma$  such that  $\mathcal{A} \in \chi_C^A[\mathcal{B}]$ . This implies  $\chi_C^A[\mathcal{A}] \subseteq \chi_C^A[\chi_C^A[\mathcal{B}]] = \chi_C^A[\mathcal{B}]$  and  $Id\chi_C^A[\mathcal{B}] \subseteq Id\chi_C^A[\mathcal{A}]$ . From  $\mathcal{B} \in \mathcal{C}Mod\Sigma$  it follows  $\chi_C^E[\Sigma] \subseteq Id\chi_C^A[\mathcal{B}]$  by (\*). Thus  $\chi_C^E[\Sigma] \subseteq Id\chi_C^A[\mathcal{A}]$  and  $\mathcal{A} \in \mathcal{C}Mod\Sigma$  by (\*). This shows that  $\chi_C^A[\mathcal{C}Mod\Sigma] \subseteq \mathcal{C}Mod\Sigma$ . ■

Because of  $R_1 \subseteq \models$ , we have  $\mathcal{C}Mod\Sigma \subseteq Mod\Sigma$  and  $\mathcal{C}IdK \subseteq IdK$  for all sets  $\Sigma \subseteq W_\tau(X)^2$  and for all  $K \subseteq Alg(\tau)$ .

The following example shows that  $R_1$  is not a Galois closed subrelation of  $\models$ . We consider the set of all identities of the greatest solid variety  $V_{HS} := Mod\{x_1(x_2x_3) \approx (x_1x_2)x_3, x_1^2 \approx x_1^4, x_1x_2x_1x_3x_1x_2x_1 \approx x_1x_2x_3x_2x_1, x_1^2x_2^2x_3 \approx x_1^2x_2x_1^2x_2x_3, x_1x_2^2x_3^2 \approx x_1x_2x_3^2x_2x_3^2\}$  of semigroups ([5]).

**Example 3.11.** Let  $\tau = (2)$  and let  $f$  be the binary operation symbol. Then there is a coloration  $C$  of  $W_{(2)}(X)$  such that  $R_1$  is not a Galois-closed subrelation of  $\models$ . We will show that there is a set  $\Sigma$  of equations with

$$\Sigma = \mathcal{C}IdV_{HS} \text{ and } V_{HS} = \mathcal{C}Mod\Sigma \not\models \Sigma = IdV_{HS}.$$

We put  $s := f(f(x, x), f(f(x, x), f(x, x)))$  and  $\alpha_s : add(s) \longrightarrow \mathbb{N}$  with  $\alpha_s(a) = 1$  for all  $a \in add(s)$ .

We set  $\Psi := (\{s \approx t \mid t \in W_{(2)}(X) \setminus \{s\}\} \cup \{t \approx s \mid t \in W_{(2)}(X) \setminus \{s\}\}) \cap IdV_{HS}$  and  $\Sigma := IdV_{HS} \setminus \Psi$ .

For  $t \in W_{(2)}(X) \setminus \{s\}$  we define  $\alpha_t : add(t) \longrightarrow \mathbb{N}$  with  $\alpha_t(a) = 0$  for all  $a \in add(t)$ .

This gives us a coloration  $C$  of  $W_{(2)}(X)$ .

If  $t \in W_{(2)}(X) \setminus \{s\}$  then for  $\rho \in Hyp(2)^\mathbb{N}$  holds  $\widehat{\rho}[t] = \widehat{\rho(0)}[t]$  by Lemma 2.5

As a consequence we have (together with Lemma 3.6)  $\chi_C^e[\Theta] = \chi^E[\Theta]$  for each set  $\Theta \subseteq W_{(2)}(X)^2$ . Since  $\chi^E$  is a closure operator, we have the idempotence property and it is easy to check that  $\chi_C^{e,n}[\Sigma] = \chi^E[\Sigma]$  for all  $n \in \mathbb{N}$ . This shows that  $\chi_C^E[\Sigma] = \chi^E[\Sigma]$ .

Hence  $\mathcal{C}Mod\Sigma = \{\mathcal{A} \mid \{\mathcal{A}\} \times \Sigma \subseteq R_1\} = \{\mathcal{A} \mid \mathcal{A} \in Alg(\tau), \chi_C^E[\Sigma] \subseteq Id\mathcal{A}\} = \{\mathcal{A} \mid \mathcal{A} \in Alg(\tau), \chi^E[\Sigma] \subseteq Id\mathcal{A}\} = V_{HS}$  since  $V_{HS}$  is defined by the hyperidentity  $f(f(x, y), z) \approx f(x, f(y, z))$ .

Further we have  $\mathcal{C}IdV_{HS} = \{r \approx t \mid V_{HS} \times \{r \approx t\} \subseteq R_1\}$   
 $= \{r \approx t \mid r \approx t \in W_\tau(X)^2, \chi_C^E[r \approx t] \subseteq IdV_{HS}\}$   
 $= \{r \approx t \mid r \approx t \in IdV_{HS}, \chi_C^E[r \approx t] \subseteq IdV_{HS}\}$   
 $= \{r \approx t \mid r \approx t \in \Sigma, \chi_C^E[r \approx t] \subseteq IdV_{HS}\} \cup \{r \approx t \mid r \approx t \in \Psi, \chi_C^E[r \approx t] \subseteq IdV_{HS}\}$   
 $= \Sigma \cup \{r \approx t \mid r \approx t \in \Psi, \chi_C^E[r \approx t] \subseteq IdV_{HS}\}$  (since  $V_{HS}$  is solid)  
 $= \Sigma \cup \emptyset$

since the multi-hypersubstitution  $\rho \in Hyp(2)^\mathbb{N}$  with  $\rho(0) = \sigma_{xy}$  and  $\rho(a) = \sigma_x$  for  $a \in \mathbb{N} \setminus \{0\}$  provides  $\widehat{\rho_C}[s] = x$  and  $\widehat{\rho_C}[t] = t$  for  $t \in W_{(2)}(X) \setminus \{s\}$ , where  $s \approx t \in IdV_{HS}$  implies  $t \neq x$ , i.e.  $\widehat{\rho_C}[s] \approx \widehat{\rho_C}[t] \notin IdV_{HS}$ .

Finally, because of  $f(x, x) \approx s \in IdV_{HS} \setminus \Sigma$  we have  $IdV_{HS} \neq \Sigma$ .

**Proposition 3.12.** *The pair  $(\chi_C^E, \chi_C^A)$  is in general not a conjugate pair of completely additive closure operators.*

**Proof:** Assume that  $(\chi_C^E, \chi_C^A)$  forms a conjugate pair of completely additive closure operators. Then for all  $\mathcal{A} \in Alg(\tau)$  and all  $s \approx t \in W_\tau(X)^2$  there holds  $s \approx t \in Id\chi_C^A[\mathcal{A}]$  iff  $\chi_C^E[s \approx t] \subseteq Id\mathcal{A}$ . In particular,  $R_1 = R_2$ . But  $R_2$  is a Galois-closed subrelation of  $\models$  and we showed in Example 3.11 that  $R_1$  is not Galois-closed. Thus  $R_1$  and  $R_2$  are different, a contradiction to  $R_1 = R_2$ .  $\blacksquare$

Since the proof of the four equivalent characterizations of  $M$ -solid varieties uses this property we cannot expect to have the same situation.

**Proposition 3.13.** *Let  $C$  be a coloration of  $W_\tau(X)$ ,  $K \subseteq \text{Alg}(\tau)$  and  $\Sigma \subseteq W_\tau(X)^2$ . Then there holds*

- (i)  $\mathcal{C}Mod\Sigma = Mod\chi_C^E[\Sigma]$ ,
- (ii)  $\mathcal{C}IdK \subseteq Id\chi_C^A[K]$ , but the converse inclusion is in general not true.

**Proof:** (i) There holds

$$\mathcal{A} \in \mathcal{C}Mod\Sigma \iff \chi_C^E[\Sigma] \subseteq Id\mathcal{A} \iff \mathcal{A} \in Mod\chi_C^E[\Sigma].$$

(ii) Let  $u \approx v \in \mathcal{C}IdK$ . Then  $\chi_C^E[u \approx v] \subseteq IdK$  and  $\chi_C^E[u \approx v] \subseteq Id\chi_C^A[K]$  by Lemma 3.9. Now we have  $u \approx v \in \chi_C^E[u \approx v] \subseteq Id\chi_C^A[K]$ . Altogether this shows that  $\mathcal{C}IdK \subseteq Id\chi_C^A[K]$ .

In order to show that the converse direction is in general not true we consider the type (2) and the coloration  $C$  of  $W_{(2)}(X)$  from Example 3.11. Moreover let  $\Sigma \subseteq W_\tau(X)^2$  be as given in Example 3.11. Then we have  $\mathcal{C}IdV_{HS} = \Sigma \neq IdV_{HS} = Id\chi^A[V_{HS}] = Id\chi_C^A[V_{HS}]$ , since we have  $V_{HS} = \chi^A[V_{HS}]$  ( $V_{HS}$  is solid) and  $\chi^A[V_{HS}] = \chi_C^A[V_{HS}]$  (Lemma 3.6). ■

**Proposition 3.14.** *Let  $C$  be a coloration of  $W_\tau(X)$ ,  $K \subseteq \text{Alg}(\tau)$  and  $\Sigma \subseteq W_\tau(X)^2$ . Then there holds:*

- (i)  $\chi_C^E[Id\mathcal{C}Mod\Sigma] \supseteq Id\mathcal{C}Mod\Sigma$ , but the converse inclusion is in general not true,
- (ii)  $\chi_C^A[Mod\mathcal{C}IdK] = Mod\mathcal{C}IdK$ .

**Proof:** (i) The inclusion is clear, since  $\chi_C^E$  is a closure operator.

In order to show that the converse direction is in general not true we consider the type (2) and the coloration  $C$  of  $W_{(2)}(X)$  from Example 3.11. Moreover let  $\Sigma \subseteq W_\tau(X)^2$  and  $s \in W_{(2)}(X)$  be as given in Example 3.11 and  $\rho \in \text{Hyp}(\tau)^\mathbb{N}$  be defined by  $\rho(0) = \sigma_{xy}$  and  $\rho(a) = \sigma_x$  for  $a \in \mathbb{N} \setminus \{0\}$ . Then we get  $s \approx f(x, x) \in IdV_{HS}$  and thus  $\widehat{\rho}_C[s] \approx \widehat{\rho}_C[f(x, x)] \in \chi_C^E[IdV_{HS}]$ , i.e.  $x \approx x^2 \in \chi_C^E[IdV_{HS}]$ . Because of  $x \approx x^2 \notin IdV_{HS}$  we obtain  $\chi_C^E[IdV_{HS}] \neq IdV_{HS}$ , i.e.  $\chi_C^E[Id\mathcal{C}Mod\Sigma] \neq Id\mathcal{C}Mod\Sigma$  since  $V_{HS} = \mathcal{C}Mod\Sigma$ .

(ii) Clearly,  $Mod\mathcal{C}IdK \subseteq \chi_C^A[Mod\mathcal{C}IdK]$ .

For the converse inclusion let  $\mathcal{A} \in \chi_C^A[Mod\mathcal{C}IdK]$ . Then there is a  $\mathcal{B} \in Mod\mathcal{C}IdK$  with  $\mathcal{A} \in \chi_C^A[\mathcal{B}]$ . We want to show that  $\mathcal{C}IdK \subseteq Id\mathcal{A}$ . For this let  $u \approx v \in \mathcal{C}IdK$ , i.e.  $\chi_C^E[u \approx v] \subseteq IdK$ . Since  $\chi_C^E$  is a closure operator we have  $\chi_C^E[\chi_C^E[u \approx v]] \subseteq IdK$  and thus  $\chi_C^E[u \approx v] \subseteq \mathcal{C}IdK$ . Since  $\mathcal{B} \in Mod\mathcal{C}IdK$  we have then  $\chi_C^E[u \approx v] \subseteq Id\mathcal{B}$ . Lemma 3.9 shows that then  $\chi_C^E[u \approx v] \subseteq Id\chi_C^A[\mathcal{B}]$ . Moreover  $\mathcal{A} \in \chi_C^A[\mathcal{B}]$  implies  $Id\chi_C^A[\mathcal{B}] \subseteq Id\mathcal{A}$ . Altogether we have  $u \approx v \in \chi_C^E[u \approx v] \subseteq Id\chi_C^A[\mathcal{B}] \subseteq Id\mathcal{A}$ . Consequently,  $\mathcal{C}IdK \subseteq Id\mathcal{A}$ .

Finally,  $\mathcal{C}IdK \subseteq Id\mathcal{A}$  means  $\mathcal{A} \in Mod\mathcal{C}IdK$ . Altogether this shows the converse inclusion  $\chi_C^A[Mod\mathcal{C}IdK] \subseteq Mod\mathcal{C}IdK$ .  $\blacksquare$

**Proposition 3.15.** *Let  $C$  be a coloration of  $W_\tau(X)$ ,  $K \subseteq Alg(\tau)$  and  $\Sigma \subseteq W_\tau(X)^2$ . Then there holds:*

- (i)  $ModId\chi_C^A[K] \subseteq CMod\mathcal{C}IdK$ , but the converse inclusion is in general not true;
- (ii)  $\mathcal{C}IdCMod\Sigma \subseteq IdMod\chi_C^E[\Sigma]$ , but the converse inclusion is in general not true.

**Proof:** (i) We have

$$\begin{aligned} ModId\chi_C^A[K] &\subseteq Mod\mathcal{C}IdK && \text{by Proposition 3.13 (ii)} \\ &= Mod\chi_C^E[\mathcal{C}IdK] && \text{by Lemma 3.10 (i)} \\ &= CMod\mathcal{C}IdK && \text{by Proposition 3.13 (i).} \end{aligned}$$

In order to show that the converse direction is in general not true we consider the type  $\tau = (2)$ . Further let  $\beta : W_\tau(X) \rightarrow \mathbb{N}$  be a bijection. (Such a bijection exists since  $W_\tau(X)$  is countable.) For  $t \in W_{(2)}(X)$  we define  $\alpha_t : add(t) \rightarrow \mathbb{N}$  with  $\alpha_t(a) = \beta(t)$  for all  $a \in add(t)$ . Then  $C := \{\alpha_t \mid t \in W_{(2)}(X)\}$  is a coloration of  $W_{(2)}(X)$ .

Let  $u \approx v \in W_\tau(X)^2$ . If  $u = v$  then obviously  $\chi_C^E[u \approx v] \subseteq \{w \approx w \mid w \in W_{(2)}(X)\} \subseteq IdV_{HS}$ . If  $u \neq v$  and  $u, v \in X$  then we get  $\chi_C^E[u \approx v] = \{u \approx v\} \not\subseteq IdV_{HS}$ . If  $u \neq v$  and  $u \notin X$  then  $\beta(u) \neq \beta(v)$  and we may consider the multi-hypersubstitution  $\rho \in Hyp(2)^\mathbb{N}$  with  $\rho(\beta(u)) = \sigma_{xy}$  and  $\rho(\beta(v)) = \sigma_x$ . Then  $\hat{\rho}_C[u] = u$  and  $\hat{\rho}_C[v] = r$ , where  $r$  is the first letter in  $v$ . Thus  $\hat{\rho}_C[u] \approx \hat{\rho}_C[v] \notin IdV_{HS}$ , i.e.  $\chi_C^E[u \approx v] \not\subseteq IdV_{HS}$ . If  $u \neq v$  and  $v \notin X$  then we get  $\chi_C^E[u \approx v] \not\subseteq IdV_{HS}$  in the dual manner. Altogether we have  $\mathcal{C}IdV_{HS} = \{w \approx w \mid w \in W_{(2)}(X)\}$  and consequently,  $CMod\mathcal{C}IdV_{HS} = Mod\{w \approx w \mid w \in W_{(2)}(X)\} = Alg(2)$  by Lemma 3.10 (i) and Proposition 3.13 (i).

On the other hand we have

$$ModId\chi_C^A[V_{HS}] = ModId\chi_C^A[V_{HS}] = ModIdV_{HS} = V_{HS}$$

by Lemma 3.6 and since  $V_{HS}$  is solid. This shows that

$$ModId\chi_C^A[V_{HS}] \neq CMod\mathcal{C}IdV_{HS}.$$

(ii) We have

$$\mathcal{C}IdCMod\Sigma \subseteq Id\chi_C^A[CMod\Sigma] = IdCMod\Sigma = IdMod\chi_C^E[\Sigma]$$

by Proposition 3.13 (ii), Lemma 3.10 (ii) and Proposition 3.13 (i).

In order to show that the converse direction is in general not true we consider the type (2) and the coloration  $C$  of  $W_{(2)}(X)$  from Example

3.11. Moreover let  $\Sigma \subseteq W_\tau(X)^2$  be defined as in Example 3.11. Then we have  $\mathcal{C}Id\mathcal{C}Mod\Sigma = \mathcal{C}IdV_{HS} = \Sigma$ .

On the other hand we have  $IdMod\chi_C^E[\Sigma] = Id\mathcal{C}Mod\Sigma = IdV_{HS} \neq \Sigma$  by Proposition 3.13 (i). Consequently,  $\mathcal{C}Id\mathcal{C}Mod\Sigma \neq IdMod\chi_C^E[\Sigma]$ . ■

#### 4. $C$ -COLORED HYPEREQUATIONAL THEORIES

**Definition 4.1.** Let  $\Sigma$  be an equational theory of type  $\tau$ . Then  $\Sigma$  is said to be a  $C$ -colored hyperequational theory if  $\mathcal{C}Mod\Sigma = Mod\Sigma$ .

$C$ -colored hyperequational theories can be characterized in the same way as usual hyperequational theories:

**Theorem 4.2.** Let  $\Sigma \in \mathcal{E}(\tau)$ . Then the following statements are equivalent:

- (i)  $\mathcal{C}Mod\Sigma = Mod\Sigma$ .
- (ii)  $\Sigma = \mathcal{C}Id\mathcal{C}Mod\Sigma$ .
- (iii)  $\Sigma = \chi_C^E[\Sigma]$ .
- (iv)  $\mathcal{C}IdMod\Sigma = \Sigma$ .

**Proof:** (i)  $\implies$  (ii): Since  $\mathcal{C}Id\mathcal{C}Mod$  is a closure operator we have  $\Sigma \subseteq \mathcal{C}Id\mathcal{C}Mod\Sigma$ . Conversely,  $\mathcal{C}Id\mathcal{C}Mod\Sigma \subseteq Id\mathcal{C}Mod\Sigma = IdMod\Sigma = \Sigma$ .

(ii)  $\implies$  (iii): Clearly,  $\Sigma \subseteq \chi_C^E[\Sigma]$ . Conversely, let  $u \approx v \in \chi_C^E[\Sigma] = \chi_C^E[\mathcal{C}Id\mathcal{C}Mod\Sigma]$  (because of (ii)). Then there is an  $s \approx t \in \mathcal{C}Id\mathcal{C}Mod\Sigma$  with  $u \approx v \in \chi_C^E[s \approx t]$ . Because of  $s \approx t \in \mathcal{C}Id\mathcal{C}Mod\Sigma$  we have  $\chi_C^E[s \approx t] \subseteq Id\mathcal{C}Mod\Sigma$ . With

$$K := \mathcal{C}Mod\Sigma \text{ and } \mathcal{C}IdK = \mathcal{C}Id\mathcal{C}Mod\Sigma = \Sigma$$

by Proposition 3.8 we get  $K = Mod\Sigma$ . Thus  $Id\mathcal{C}Mod\Sigma = IdMod\Sigma = \Sigma$  since  $\Sigma \in \mathcal{E}(\tau)$ . Altogether we have

$$u \approx v \in \chi_C^E[s \approx t] \subseteq Id\mathcal{C}Mod\Sigma = \Sigma.$$

This shows that  $\chi_C^E[\Sigma] \subseteq \Sigma$ .

(iii)  $\implies$  (i):  $\mathcal{C}Mod\Sigma \subseteq Mod\Sigma$  is clear. Conversely, let  $\mathcal{A} \in Mod\Sigma$ . Then  $\Sigma \subseteq Id\mathcal{A}$ . Because of (iii) we have then  $\chi_C^E[\Sigma] \subseteq Id\mathcal{A}$ , i.e.  $\mathcal{A} \in \mathcal{C}Mod\Sigma$ . This shows that  $Mod\Sigma \subseteq \mathcal{C}Mod\Sigma$ .

(i)  $\Leftrightarrow$  (iv): Suppose that  $\mathcal{C}IdMod\Sigma = \Sigma$ . Then we put

$$K := \mathcal{C}Mod\mathcal{C}IdMod\Sigma.$$

Now we have

$$\mathcal{C}Mod\Sigma = \mathcal{C}Mod\mathcal{C}IdMod\Sigma = K$$

and

$$\mathcal{C}IdK = \mathcal{C}Id\mathcal{C}Mod\mathcal{C}IdMod\Sigma = \mathcal{C}IdMod\Sigma$$

(since  $(CId, CMod)$  is a Galois-connection). By Proposition 3.8 we get  $K = Mod\Sigma$ , i.e.  $CModCIdMod\Sigma = Mod\Sigma$ ,  $CMod\Sigma = Mod\Sigma$  and  $\Sigma$  is a  $C$ -colored hyperequational theory.

Suppose that  $\Sigma$  is a  $C$ -colored hyperequational theory. We use that  $\Sigma = IdMod\Sigma \supseteq CIdMod\Sigma$ . Now let  $u \approx v \in \Sigma$ . Assume that  $u \approx v \notin CIdMod\Sigma$ . Then  $\chi_C^E[u \approx v] \not\subseteq IdMod\Sigma$  and in particular there is an  $\mathcal{A} \in Mod\Sigma$  with  $\chi_C^E[u \approx v] \not\subseteq Id\mathcal{A}$ . Since  $\Sigma$  is a  $C$ -colored hyperequational theory we have  $Mod\Sigma = CMod\Sigma$ , i.e.  $\mathcal{A} \in CMod\Sigma$  and  $\chi_C^E[\Sigma] \subseteq Id\mathcal{A}$ . Because of  $u \approx v \in \Sigma$  we have  $\chi_C^E[u \approx v] \subseteq \chi_C^E[\Sigma]$  and thus  $\chi_C^E[u \approx v] \subseteq Id\mathcal{A}$ , a contradiction. Consequently,  $u \approx v \in CIdMod\Sigma$ . This shows that  $\Sigma \subseteq CIdMod\Sigma$ . ■

When  $\Sigma$  is a  $C$ -colored hyperequational theory, then  $Mod\Sigma = \chi_C^A[Mod\Sigma]$ . To see this, suppose that  $Mod\Sigma \neq \chi_C^A[Mod\Sigma]$ , then  $Mod\Sigma \neq \chi^A[Mod\Sigma]$  by Lemma 3.6, i.e.  $Mod\Sigma$  is not solid. Then there is an  $\mathcal{A} \in Mod\Sigma$  with  $\chi^E[\Sigma] \not\subseteq Id\mathcal{A}$ . Because of  $\chi^E[\Sigma] \subseteq \chi_C^E[\Sigma]$  we have  $\chi_C^E[\Sigma] \not\subseteq Id\mathcal{A}$ , i.e.  $\mathcal{A} \notin CMod\Sigma$  contradicts  $CMod\Sigma = Mod\Sigma$ .

The following example shows that the converse is not true, i.e.  $C$ -colored hyperequational theories cannot be characterized by the condition  $Mod\Sigma = \chi_C^E[Mod\Sigma]$ .

**Example 4.3.** Let  $\tau = (2)$ . We use the coloration  $C$  from Example 3.11. Further we take  $\Sigma := IdV_{HS}$ . Then we have  $ModIdV_{HS} = V_{HS} = \chi^A[V_{HS}]$ , since  $V_{HS}$  is solid. By Lemma 3.6 we have  $V_{HS} = \chi_C^A[V_{HS}] = \chi_C^A[ModIdV_{HS}]$ .

Now we show that  $CModIdV_{HS} \neq V_{HS} = ModIdV_{HS}$ . Actually we will show that  $CModIdV_{HS} \subseteq \mathbf{B}$ , where  $\mathbf{B}$  denotes the variety of bands. We consider the term  $s$  and the hypersubstitution  $\rho$  from Example 3.11 and note that the application of  $\rho$  to the identity  $s \approx f(x, x) \in IdV_{HS}$  provides the idempotent law. Thus  $x \approx x^2 \in \chi_C^E[IdV_{HS}]$ . Let  $\mathcal{A} \in CModIdV_{HS}$ . Then  $\chi_C^E[IdV_{HS}] \subseteq Id\mathcal{A}$ , and in particular  $x \approx x^2 \in Id\mathcal{A}$ , i.e.  $\mathcal{A} \in \mathbf{B}$ .

## 5. CHARACTERIZATIONS OF COLORED SOLID VARIETIES

In Section 3 colored solid varieties  $V$  were defined by the property  $IdV = \chi_C^E[IdV]$ . We get the following characterization:

**Theorem 5.1.** *Let  $C$  be a coloration of  $W_\tau(X)$  and  $V$  be a variety of type  $\tau$ . Then the following statements (i)-(iii) are equivalent:*

- (i)  $CModIdV = V$ .
- (ii)  $IdV = CIdV$ .



(iii)  $\chi_C^E[IdV] = IdV$ .

Further, each of the statements (i)-(iii) implies both  $V = \mathcal{C}Mod\mathcal{C}IdV$  and  $\chi_C^A[V] = V$ , but the converse implications are in general not true.

**Proof:** (i) $\Rightarrow$ (ii) Suppose that  $\mathcal{C}ModIdV = V$ . Then  $IdV$  is a  $C$ -colored hyperequational theory, i.e.  $ModIdV = \mathcal{C}ModIdV$  and  $\mathcal{C}Id\mathcal{C}ModIdV = IdV$ . Thus  $\mathcal{C}IdV = IdV$ , i.e.  $V$  is  $C$ -colored solid.

(ii) $\Rightarrow$ (iii) Since  $\chi_C^E$  is a closure operator we have  $IdV \subseteq \chi_C^E[IdV]$ . Conversely, let  $u \approx v \in \chi_C^E[IdV]$ . Then there is an  $s \approx t \in IdV$  with  $u \approx v \in \chi_C^E[s \approx t]$ . From  $s \approx t \in IdV$  it follows that  $s \approx t \in \mathcal{C}IdV$  by (ii), i.e.  $\chi_C^E[s \approx t] \subseteq IdV$ . Altogether we have  $u \approx v \in IdV$ . This shows that  $\chi_C^E[IdV] \subseteq IdV$ .

(iii)  $\Rightarrow$ (i)  $IdV$  is an equational theory. Thus we can use Theorem 4.2 and from  $IdV = \chi_C^E[IdV]$  it follows that  $\mathcal{C}ModIdV = ModIdV$  and  $IdV = \mathcal{C}Id\mathcal{C}ModIdV$ . This gives

$$IdV = \mathcal{C}Id\mathcal{C}ModIdV = \mathcal{C}IdModIdV = \mathcal{C}IdV.$$

Moreover we have

$$V \subseteq \mathcal{C}Mod\mathcal{C}IdV = \mathcal{C}ModIdV = ModIdV = V.$$

This shows that  $V = \mathcal{C}ModIdV$ .

Suppose that  $IdV = \mathcal{C}IdV$ . We show that  $V = \mathcal{C}Mod\mathcal{C}IdV$ . First we have that  $V \subseteq \mathcal{C}Mod\mathcal{C}IdV$  since  $\mathcal{C}Mod\mathcal{C}Id$  is a closure operator. Conversely we get that  $\mathcal{C}ModIdV \subseteq ModIdV = V$ . Using  $IdV = \mathcal{C}IdV$  one obtains  $\mathcal{C}Mod\mathcal{C}IdV \subseteq V$ .

Now we show that  $V = \chi_C^A[V]$ . Assume that  $V \neq \chi_C^A[V]$ . Then  $V \neq \chi^A[V]$  by Lemma 3.6, i.e.  $V$  is not solid and  $\chi^E[IdV] \not\subseteq IdV$ . Because of  $\chi^E[IdV] \subseteq \chi_C^E[IdV]$  we have that  $\chi_C^E[IdV] \not\subseteq IdV$ . This shows that  $IdV \not\subseteq \mathcal{C}IdV$ , contradicting  $IdV = \mathcal{C}IdV$ .

Finally we prove that the opposite implication is not satisfied. Let  $\tau = (2)$ . Then there are a coloration  $C$  of  $W_{(2)}(X)$ , a variety  $V$  of type (2) with  $V = \mathcal{C}Mod\mathcal{C}IdV$  and  $V = \chi_C^A[V]$  such that  $\mathcal{C}IdV \neq IdV$ . Indeed, we take the coloration  $C$  and the set  $\Sigma$  from the proof of Example 3.11. There we have shown that  $\mathcal{C}Mod\Sigma = V_{HS}$  and  $\mathcal{C}IdV_{HS} = \Sigma$ . So we have  $V_{HS} = \mathcal{C}Mod\Sigma = \mathcal{C}Mod\mathcal{C}IdV_{HS}$ . On the other hand there holds  $V_{HS} = \chi^A[V_{HS}]$  since  $V_{HS}$  is solid. Thus  $V_{HS} = \chi_C^A[V_{HS}]$  by Lemma 3.6. Moreover there holds  $\mathcal{C}IdV_{HS} = \Sigma \neq IdV_{HS}$ . ■

$C$ -colored hyperequational theories also have the following properties.

**Proposition 5.2.** *Let  $C$  be a coloration of  $W_\tau(X)$  and  $\Sigma$  be an equational theory. If  $Id\mathcal{C}Mod\Sigma = \Sigma$  then  $\Sigma$  is a  $C$ -colored hyperequational theory. The converse direction is not true.*

**Proof:** Suppose that  $Id\mathcal{C}Mod\Sigma = \Sigma$ . Clearly,  $\Sigma \subseteq \mathcal{C}Id\mathcal{C}Mod\Sigma$ . Let  $u \approx v \in \mathcal{C}Id\mathcal{C}Mod\Sigma$ . Then  $\chi_C^E[u \approx v] \subseteq Id\mathcal{C}Mod\Sigma$ , i.e.  $u \approx v \in \chi_C^E[u \approx v] \subseteq Id\mathcal{C}Mod\Sigma = \Sigma$  and  $u \approx v \in \Sigma$ . This shows  $\mathcal{C}Id\mathcal{C}Mod\Sigma \subseteq \Sigma$ . Altogether we have  $\Sigma = \mathcal{C}Id\mathcal{C}Mod\Sigma$ . By Theorem 4.2 we have  $\mathcal{C}Mod\Sigma = Mod\Sigma$ , i.e.  $\Sigma$  is a  $C$ -colored hyperequational theory.

In order to show that the converse direction is not true we consider the type (2) and the coloration  $C$  of  $W_{(2)}(X)$  from Example 3.11. Moreover let  $\Sigma \subseteq W_\tau(X)^2$  be defined as in Example 3.11. Then we have  $\Sigma = \mathcal{C}IdV_{HS} = \mathcal{C}Id\mathcal{C}Mod\Sigma$ , i.e.  $\Sigma$  is a  $C$ -colored hyperequational theory. But there holds  $Id\mathcal{C}Mod\Sigma = IdV_{HS} \neq \Sigma$ . ■

**Proposition 5.3.** *Let  $C$  be a coloration of  $W_\tau(X)$  and  $K$  be a class of algebras of type  $\tau$ . Then  $Mod\mathcal{C}IdK = \mathcal{C}Mod\mathcal{C}IdK$ .*

**Proof:** We have  $\mathcal{C}Mod\mathcal{C}IdK = \{\mathcal{A} \mid \mathcal{A} \in Alg(\tau), \chi_C^E[\mathcal{C}IdK] \subseteq Id\mathcal{A}\} = \{\mathcal{A} \mid \mathcal{A} \in Alg(\tau), \mathcal{C}IdK \subseteq Id\mathcal{A}\} = Mod\mathcal{C}IdK$  by Lemma 3.10 (i). ■

## 6. EXAMPLES

For a variety to be  $C$ -colored solid, it must satisfy all identities obtained by applying all multi-hypersubstitutions to all identities of the variety. This can be difficult to verify. In the hyperidentity case, if we want to check whether a variety of the form  $V = Mod\Sigma$  is solid, we have to apply the hypersubstitutions only to the set  $\Sigma$ . This is based on the following theorem:

**Theorem 6.1.** [1] *Let  $K \subseteq Alg(\tau)$  and  $\Sigma \subseteq W_\tau(X)^2$  and let  $\mathcal{M} \subseteq Hyp(\tau)$  be a submonoid. Then the following holds:*

- (i)  $\chi_M^E[\Sigma] \subseteq IdMod\Sigma \iff IdMod\Sigma = H_M IdH_M Mod\Sigma$ .
- (ii)  $\chi_M^E[\Sigma] \subseteq IdMod\Sigma \iff \chi_M^E[IdMod\Sigma] \subseteq IdMod\Sigma$ .
- (iii)  $\chi_M^A[K] \subseteq ModIdK \iff ModIdK = H_M ModH_M IdK$ .
- (iv)  $\chi_M^A[K] \subseteq ModIdK \iff \chi_M^A[ModIdK] \subseteq ModIdK$ .

For multi-hypersubstitutions and colored terms we have:

**Theorem 6.2.** *Let  $K \subseteq Alg(\tau)$ , let  $C$  be a coloration of  $W_\tau(X)$  and  $\Sigma \subseteq W_\tau(X)^2$ . Then the following holds:*

- (i)  $\chi_C^E[\Sigma] \subseteq IdMod\Sigma \iff IdMod\Sigma \supseteq \mathcal{C}Id\mathcal{C}Mod\Sigma$ .
- (ii)  $\chi_C^E[\Sigma] \subseteq IdMod\Sigma \iff \chi_C^E[IdMod\Sigma] \subseteq IdMod\Sigma$ .
- (iii)  $\chi_C^A[K] \subseteq ModIdK \iff ModIdK = \mathcal{C}Mod\mathcal{C}IdK$ .

$$(iv) \chi_C^A[K] \subseteq ModIdK \iff \chi_C^A[ModIdK] \subseteq ModIdK.$$

The converse implications are in general not satisfied.

**Proof:** (i): Suppose that  $\chi_C^E[\Sigma] \subseteq IdMod\Sigma$ . Then we have

$$\mathcal{C}Id\mathcal{C}Mod\Sigma \subseteq IdMod\chi_C^E[\Sigma] \subseteq IdModIdMod\Sigma = IdMod\Sigma$$

by Proposition 3.15 (i).

Suppose that  $IdMod\Sigma \supseteq \mathcal{C}Id\mathcal{C}Mod\Sigma$ . Then we have

$$\chi_C^E[\Sigma] \subseteq \mathcal{C}Id\mathcal{C}Mod\chi_C^E[\Sigma] = \mathcal{C}Id\mathcal{C}Mod\Sigma \subseteq IdMod\Sigma$$

because of  $\mathcal{C}Mod\Sigma = Mod\chi_C^E[\Sigma] = Mod\chi_C^E[\chi_C^E[\Sigma]] = \mathcal{C}Mod\chi_C^E[\Sigma]$  (Proposition 3.13).

In order to show that the implication

$$\chi_C^E[\Sigma] \subseteq IdMod\Sigma \implies IdMod\Sigma = \mathcal{C}Id\mathcal{C}Mod\Sigma$$

is in general not true we consider the type (2) and the coloration  $C$  of  $W_{(2)}(X)$  from Example 3.11. Moreover let  $\Sigma \subseteq W_\tau(X)^2$  be defined as in Example 3.11. Then we have

$$\chi_C^E[\chi_C^E[\Sigma]] = \chi_C^E[\Sigma] \subseteq IdMod\chi_C^E[\Sigma].$$

On the other hand we have

$$\mathcal{C}Id\mathcal{C}Mod\chi_C^E[\Sigma] = \mathcal{C}Id\mathcal{C}Mod\Sigma = \mathcal{C}IdV_{HS} = \Sigma$$

and

$$IdMod\chi_C^E[\Sigma] = Id\mathcal{C}Mod\Sigma = IdV_{HS} \neq \Sigma.$$

This shows that indeed  $\chi_C^E[\chi_C^E[\Sigma]] \subseteq IdMod\chi_C^E[\Sigma]$  but  $IdMod\chi_C^E[\Sigma] \neq \mathcal{C}Id\mathcal{C}Mod\chi_C^E[\Sigma]$ .

(ii): This implication is clear since  $\chi_C^E[\Sigma] \subseteq \chi_C^E[IdMod\Sigma]$ .

In order to show that the converse direction is in general not true we consider the type (2) and the coloration  $C$  of  $W_{(2)}(X)^2$  from Example 3.11. Moreover let  $\Sigma \subseteq W_\tau(X)$  be defined as in Example 3.11. Then we have  $\chi_C^E[\chi_C^E[\Sigma]] = \chi_C^E[\Sigma] \subseteq IdMod\chi_C^E[\Sigma]$ .

On the other hand we have  $IdMod\chi_C^E[\Sigma] = IdV_{HS}$ . Further we have  $x \approx x^2 \in \chi_C^E[IdV_{HS}]$ . Since  $x \approx x^2 \notin IdV_{HS}$  we have  $\chi_C^E[IdV_{HS}] \not\subseteq IdV_{HS}$  and thus

$$\chi_C^E[IdMod\chi_C^E[\Sigma]] = \chi_C^E[IdV_{HS}] \not\subseteq IdV_{HS} = IdMod\chi_C^E[\Sigma].$$

(iii): Suppose that  $ModIdK = \mathcal{C}Mod\mathcal{C}IdK$ . Then we have  $\chi_C^A[K] \subseteq ModId\chi_C^A[K] \subseteq \mathcal{C}Mod\mathcal{C}IdK = ModIdK$  by Proposition 3.15 (i).

In order to show that the converse implication is in general not true we consider the type (2) and the coloration  $C$  of  $W_{(2)}(X)$  from Example 3.11. Then we have  $\chi_C^A[V_{HS}] = V_{HS} \subseteq ModIdV_{HS}$ .

On the other hand we have

$$\begin{aligned} \mathcal{C}Mod\mathcal{C}IdV_{HS} &= Mod\mathcal{C}IdV_{HS} \\ &= Mod\{w \approx w \mid w \in W_{(2)}(X)\} = Alg(2) \neq V_{HS} = ModIdV_{HS} \end{aligned}$$

(see proof of Proposition 3.15 (i)).  
(iv): is clear. ■

The following example shows once more that  $(\chi_C^E, \chi_C^A)$  does not form a conjugate pair.

**Proposition 6.3.** *If  $\tau = (n_i)_{i \in I}$  with  $n_k \geq 2$  for some  $k \in I$  then there are a coloration  $C$  of  $W_\tau(X)$ , terms  $s, t \in W_\tau(X)$ , and a multi-hypersubstitution  $\rho \in Hyp(\tau)^\mathbb{N}$  and an algebra  $\mathcal{A}$  of type  $\tau$  such that*

$$\mathcal{A} \models \widehat{\rho}_C[s] \approx \widehat{\rho}_C[t] \iff \rho[\mathcal{A}] \models s \approx t$$

*is not valid.*

**Proof:** We will show that for the free algebra  $\mathcal{F}_\tau(X)$  of type  $\tau$  over an alphabet  $X$  there are a coloration  $C$  of  $W_\tau(X)$  and some  $s, t \in W_\tau(X)$  and an  $\rho \in Hyp(\tau)^\mathbb{N}$  such that  $\widehat{\rho}[s] \approx \widehat{\rho}[t] \in Id\mathcal{F}_\tau(X)$  but  $s \approx t \notin Id\rho[\mathcal{F}_\tau(X)]$ .

Let  $k \in I$  with  $n_k \geq 2$ . We put  $s := f_k(f_k(x_1, \dots, x_1), x_2, \dots, x_2)$  and  $t := f_k(f_k(x_1, \dots, x_1), x_1, \dots, x_1)$  and we consider a coloration  $C$  of  $W_\tau(X)$  with the following properties:

- (i)  $\alpha_s(j) = 0$  for all  $j \in add(s)$ ,
- (ii)  $\alpha_t(j) = 0$  for all  $j \in add(t)$ ,
- (iii)  $\alpha_{f_k(x_1, \dots, x_{n_k})}(j) = 1$  for all  $j \in add(f_k(x_1, \dots, x_{n_k}))$ .

Finally, let  $\rho \in Hyp^\mathbb{N}$  given by  $\rho(0) = \sigma_{x_1}$  and  $\rho(j) = \tilde{\sigma}$  for  $j \in \mathbb{N} \setminus \{0\}$ , where  $\sigma_{x_1}$  is defined by  $\sigma_{x_1} : f_j \longrightarrow x_1$  for all  $j \in I$  and  $\tilde{\sigma}$  is defined by  $\tilde{\sigma} : f_j \longrightarrow x_{n_j}$  for all  $j \in I$ .

Then we have  $\widehat{\rho}_C[s] = \widehat{\rho}_C[t] = x_1$ . This shows that  $\widehat{\rho}_C[s] \approx \widehat{\rho}_C[t] \in Id\mathcal{F}_\tau(X)$ .

On the other hand we have

$$f_k^{\rho[\mathcal{F}_\tau(X)]} = \widehat{\rho}_C[f_k(x_1, \dots, x_{n_k})]^{\mathcal{F}_\tau(X)} = x_{n_k}^{\mathcal{F}_\tau(X)}.$$

Then the following holds:

$$\begin{aligned} s^{\rho[\mathcal{F}_\tau(X)]}(x_1, x_2) &= f_k^{\rho[\mathcal{F}_\tau(X)]}(f_k^{\rho[\mathcal{F}_\tau(X)]}(x_1, \dots, x_1), x_2, \dots, x_2) \\ &= x_{n_k}^{F_\tau(X)}(x_{n_k}^{\mathcal{F}_\tau(X)}(x_1, \dots, x_1), x_2, \dots, x_2) \\ &= x_2 \end{aligned}$$

and

$$\begin{aligned}
t^{\rho[\mathcal{F}_\tau(X)]}(x_1, x_2) &= f_k^{\rho[\mathcal{F}_\tau(X)]}(f_k^{\rho[\mathcal{F}_\tau(X)]}(x_1, \dots, x_1), x_1, \dots, x_1) \\
&= x_{n_k}^{\mathcal{F}_\tau(X)}(x_{n_k}^{\mathcal{F}_\tau(X)}(x_1, \dots, x_1), x_1, \dots, x_1) \\
&= x_1.
\end{aligned}$$

This shows  $s^{\rho[\mathcal{F}_\tau(X)]} \neq t^{\rho[\mathcal{F}_\tau(X)]}$  and  $s \approx t \notin Id\rho[\mathcal{F}_\tau(X)]$ .  $\blacksquare$

**Example 6.4.** In the next example we will determine a collection of colorations  $C$  of  $W_{(2)}(X)$  such that all solid varieties of bands are  $C$ -colored solid. There are exactly three nontrivial solid varieties of bands: the variety  $RB$  of all rectangular bands, the variety  $NB$  of all normal bands, and the variety  $RegB$  of all regular bands (see [4]).

We split the set  $W_{(2)}(X)$  into two sets  $A$  and  $B$ :

$$\begin{aligned}
A &:= \bigcup \{W_{(2)}(\{w\}) \mid w \in X\} \text{ and} \\
B &:= W_{(2)}(X) \setminus A.
\end{aligned}$$

By  $\mathcal{F}$  we denote the set of all mappings

$$\beta : \bigcup \{ \{(t, a) \mid a \in \text{add}(t)\} \mid t \in A \} \rightarrow \mathbb{N}.$$

For  $\beta \in \mathcal{F}$  we define a coloration  $C_\beta = \{\alpha_t^\beta \mid t \in W_{(2)}(X)\}$  as follows: For  $t \in W_{(2)}(X)$  and  $a \in \text{add}(t)$  we put

$$\begin{aligned}
\alpha_t^\beta(a) &= 1 && \text{if } t \in B; \\
\alpha_t^\beta(a) &= \beta(t, a) && \text{if } t \in A.
\end{aligned}$$

Then the varieties  $RB$ ,  $NB$ , and  $RegB$  are  $C_\beta$ -colored solid. Indeed, let  $V$  be one of the varieties  $RB$ ,  $NB$ , and  $RegB$  and let  $s \approx t \in IdV$ . Further let  $\rho \in Hyp(2)^\mathbb{N}$ .

If  $s, t \in A$  then there is a variable  $w \in X$  such that  $s, t \in W_{(2)}(\{w\})$ . (Otherwise the identity  $s \approx t$  provides the identity  $x \approx y$  because of the associative and the idempotent law.) Thus  $\widehat{\rho}_{C_\beta}[s], \widehat{\rho}_{C_\beta}[t] \in W_{(2)}(\{w\})$ . Because of the associative and the idempotent law  $\widehat{\rho}_{C_\beta}[s] \approx \widehat{\rho}_{C_\beta}[t]$  is an identity in  $V$ .

If  $s, t \in B$  then  $\widehat{\rho}_{C_\beta}[s] = \widehat{\rho(1)}[s]$  and  $\widehat{\rho}_{C_\beta}[t] = \widehat{\rho(1)}[t]$  by Lemma 2.5. Since  $V$  is solid, we have  $\widehat{\rho(1)}[s] \approx \widehat{\rho(1)}[t] \in IdV$  and thus  $\widehat{\rho}_{C_\beta}[s] \approx \widehat{\rho}_{C_\beta}[t] \in IdV$ .

If  $s \in A$  and  $t \in B$  then there is a  $w \in X$  such that  $s \in W_{(2)}(\{w\})$ . Since  $V$  is a variety of bands we have  $\widehat{\rho}_{C_\beta}[s] \approx w \approx \widehat{\rho(1)}[s]$ . By Lemma

2.5 we have  $\widehat{\rho}_{C_\beta}[t] = \widehat{\rho(1)}[t]$ , Since  $V$  is solid we have  $\widehat{\rho(1)}[s] \approx \widehat{\rho(1)}[t] \in IdV$  and thus  $\widehat{\rho}_{C_\beta}[s] \approx \widehat{\rho}_{C_\beta}[t] \in IdV$ .

If  $s \in B$  and  $t \in A$  then we get in a similar way that  $\widehat{\rho}_{C_\beta}[s] \approx \widehat{\rho}_{C_\beta}[t] \in IdV$ .

**Example 6.5.** We consider the following coloration  $C = \{\alpha_t \mid t \in W_{(2)}(X)\}$  of  $W_{(2)}(X)$ : for  $t \in W_{(2)}(X)$  and  $a \in add(t)$  we put

$$\begin{aligned} \alpha_t(a) &= 1 \quad \text{if } t = f(x, x); \\ \alpha_t(a) &= 2 \quad \text{if } t \neq f(x, x). \end{aligned}$$

The varieties  $RB$ ,  $NB$ , and  $RegB$  are the only nontrivial  $C$ -colored solid varieties of semigroups. Indeed, by Example 6.4,  $RB$ ,  $NB$ , and  $RegB$  are  $C$ -colored solid.

Conversely, let  $V$  be a  $C$ -colored solid variety of semigroups and  $\rho \in Hyp(2)^{\mathbb{N}}$  with

$$\rho(1) = \sigma_x \text{ and } \rho(i) = \sigma_{xy} \text{ for } 2 \leq i \in \mathbb{N}.$$

Since  $V$  is  $C$ -colored solid,  $V$  is solid and thus  $V \subseteq V_{HS}$ , i.e.  $f(x, x) \approx f(f(x, x), f(x, x)) \in IdV$ . Then we have:

$$\begin{aligned} x &\approx \widehat{\rho(1)}[f(x, x)] \approx \widehat{\rho_C}[f(x, x)] \text{ (by Lemma 2.5)} \\ &\approx \widehat{\rho_C}[f(f(x, x), f(x, x))] \text{ (} V \text{ is } C\text{-colored solid)} \\ &\approx \widehat{\rho(2)}[f(f(x, x), f(x, x))] \text{ (by Lemma 2.5)} \\ &\approx f(f(x, x), f(x, x)) \\ &\approx f(x, x). \end{aligned}$$

This shows that  $V$  is a variety of bands. But  $RB$ ,  $NB$ , and  $RegB$  are the only nontrivial solid varieties of bands.

## REFERENCES

- [1] K. Denecke, M. Reichel, *Monoids of hypersubstitutions and M-solid varieties*, Contributions to General Algebra 9, Verlag B.G. Teubner, Stuttgart, 1995, 117-125.
- [2] K. Denecke, S. L. Wismath, *Hyperidentities and Clones*, Gordon and Breach Science Publishers, 2000.
- [3] K. Denecke, S. L. Wismath, *Universal Algebra and Applications in Theoretical Computer Science*, Chapman and Hall/ CRC, 2002.
- [4] E. Graczyńska, *On normal and regular identities and hyperidentities*, Universal and Applied Algebra, Turawa, Poland 3 - 7 May 1988, World Scientific (1989), 107-135.

- [5] L. Polák, *On hyperassociativity*, Algebra Universalis **36** No. 3 (1996), 363-378.
- [6] L. Polák, *All solid varieties of semigroups*, J. Algebra **219** No. 2 (1999), 421-436.

Authors' Addresses:

K. Denecke  
 Universität Potsdam  
 Fachbereich Mathematik  
 Postfach 601553  
 D-14415 Potsdam  
 email:kdenecke@rz.uni-potsdam.de

J. Koppitz  
 Universität Potsdam  
 Fachbereich Mathematik  
 Postfach 601553  
 D-14415 Potsdam  
 email:koppitz@rz.uni-potsdam.de

Sl. Shtrakov  
 South-West-University Blagoevgrad  
 Faculty of Mathematics and Natural Sciences  
 2700 Blagoevgrad, Bulgaria  
 e-mail:shtrakov@swu.bg